

Erratum to: Plant pattern-recognition receptors controlling innate immunity

Lei Li, Yufei Yu, Zhaoyang Zhou^{*} & Jian-Min Zhou^{**}

State Key Laboratory of Plant Genomics, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China

Erratum to: SCIENCE CHINA Life Sciences, September 2016 Vol.59 No.9: 878–888

doi: 10.1007/s11427-016-0115-2

The abstract and keywords are missing in the original version.

Abstract

Plants are exposed to numerous potential pathogenic microbes. To counter the threat, plants have evolved diverse pattern-recognition receptors (PRRs), which are receptor kinases (RKs) and receptor proteins (RPs) specialized to detect conserved pathogen/microbe-associated molecular patterns (PAMPs/MAMPs). Although only a handful of RKs and RPs are known PRRs, they belong to the receptor-like kinase (RLK) and receptor-like protein (RLP) superfamilies that undergo lineage-specific expansion, suggesting that many of these RLKs and RLPs are potential PRRs. Analyses of existing PRRs have uncovered ligand-induced RLK-RK or RLK-RP oligomerization as a common mechanism for immune activation. PRRs can recruit additional components to form dynamic receptor complexes, which mediate specific cellular responses. Detailed analyses of these components are shedding light on molecular mechanisms underlying the regulation of PRR activity and downstream signaling.

Keywords

receptor kinase, receptor protein, receptor-like cytoplasmic kinase, plant innate immunity

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s11427-016-0115-2>

^{*}Corresponding author (email: zhaoyangzhou@genetics.ac.cn)

^{**}Corresponding author (email: jmzhou@genetics.ac.cn)